

NAME: Patient nty2T59
ACC #: 48162 **DOB:** 1/1/1900
SEX:
SPECIMEN TYPE:
COLLECTION DATE: 1/1/1900
RECEIVED DATE: 1/1/1900
REPORT DATE: 2/8/2018

FOR ACADEMIC PURPOSES ONLY - NOT FOR CLINICAL USE

Test Details

Gene	Genotype	Phenotype	Alleles Tested
CYP2C19	*1/*2	Intermediate Metabolizer	*2, *3, *4, *4B, *5, *6, *7, *8, *9, *17
CYP2D6	*2/*35	Normal Metabolizer	*2, *3, *4, *4M, *6, *7, *8, *9, *10, *12, *14A, *14B, *17, *29, *35, *41
CYP2C9	*1/*11	Intermediate Metabolizer	*2, *3, *4, *5, *6, *11
VKORC1	-1639G>A G/A	Intermediate Warfarin Sensitivity	-1639G>A
Apolipoprotein E	Indeterminate	Unknown Phenotype	ε2, ε4, (ε3 is reference)
CYP3A5	*3/*3/*8	Poor Metabolizer	*1D, *2, *3, *3B, *3C, *6, *7, *8, *9
CYP1A2	*1A/*1A	Normal Metabolizer- Possible Inducibility	*1C, *1D, *1F, *1K, *1L, *1V, *1W
SLCO1B1	521T>C T/T	Normal Function	521T>C, 388A>G
COMT	Val158Met G/G	High/Normal COMT Activity	Val158Met
OPRM1	A118G A/A	Normal OPRM1 Function	A118G
CYP3A4	*1/*22	Intermediate Metabolizer	*1B, *2, *3, *12, *17, *22
CYP2B6	Indeterminate	Unknown Phenotype	*6, *9
MTHFR	1298A>C AA 677C>T TT	Increased Risk of Hyperhomocysteinemia	1298A>C, 677C>T
MTHFR	677C>T TT	Reduced MTHFR Activity	1298A>C, 677C>T
Factor II	20210G>A GG	No Increased Risk of Thrombosis	20210G>A, 1691G>A
Factor V Leiden	1691G>A GG		

Additional Test Results (added to this original report)

HLA-B*15:02	negative/negative	Negative	HLA-A*31:01	negative/negative	Negative
HLA-B*57:01	negative/negative	Negative			
HLA-B*58:01	negative/negative	Negative			

Limitation: This test will not detect all the known alleles that result in altered or inactive tested genes. This test does not account for all individual variations in the individual tested. Absence of a detectable gene mutation does not rule out the possibility that a patient has different phenotypes due to the presence of an undetected polymorphism or due to other factors such as drug-drug interactions, comorbidities and lifestyle habits.

Methodology: Array based assays detect listed alleles, including all common and most rare variants with known clinical significance at analytical sensitivity and specificity >99%.

Disclaimer: Manchester University developed the Genotype test. The performance characteristics of this test were determined by Manchester University. It has not been cleared or approved by the U.S. Food and Drug Administration.

Translational Software Disclaimer: The information presented on this report is provided as general educational health information. The content is not intended to be a substitute for professional medical advice, diagnosis, or treatment. Only a physician, pharmacist or other healthcare professional should advise a patient on the use of the medications prescribed.

The pharmacogenetic assay involves use of reporting software and genotype-phenotype associations performed by Translational Software (www.translationalsoftware.com). The software has not been evaluated by the Food and Drug Administration. The software, and the report generated by the software, is not intended to diagnose, treat, cure, or prevent any disease. A qualified designee within the lab uses Translational Software to generate and subsequently review the report. The pharmacogenetic report is one of multiple pieces of information that clinicians should consider in guiding their therapeutic choice for each patient. It remains the responsibility of the health-care provider to determine the best course of treatment for a patient. Adherence to dose guidelines does not necessarily assure a successful medical outcome.